

AGPRO BIO-Safe

A fast acting coconut oil derivative weedkiller for the non-selective control of a wide range of annual and some perennial weeds.

Active Ingredient: 700g/litre fatty acids in the form of an emulsifiable concentrate.

GENERAL INSTRUCTIONS:

AGPRO BIO-Safe works only on direct contact with foliage, causing rapid wilting of the leaves. The effect will be noticed within hours. It is most effective on actively growing weeds and when applied in hot, sunny conditions.

NZNO Approval number:
Batch No:
Net Contents: 20 Litres
Registered pursuant to the
ACVM Act 1997 No. 5734
See www.nzfsa.govt.nz/acvm
for registration conditions

Registered to and Distributed by:
AGPRO NZ LTD
10 Polaris Place,
East Tamaki.
P.O.Box 58-963
East Tamaki,
AUCKLAND.
PH 0-9-273 3456
FAX 0-9-273 3457
WEB www.agpro.co.nz

HAZARD CLASSIFICATIONS: 6.3A, 8.3A, 9.1C.

FIRST AID:

- If swallowed give water to dilute. Do not induce vomiting.
- For advice contact National Poisons Centre (0800 POISON 764-766) or a doctor immediately.
- If skin or hair contact occurs, remove contaminated clothing and flush skin and hair with running water.
- If splashed in eyes, wash out immediately with water.

PRECAUTIONS:

Store in the original unopened container in a cool dry place, avoiding sunlight and away from foodstuffs and drink containers.
Do not get into eyes or on skin. In case of contact, flush skin or eyes with plenty of water for 15 minutes; for eyes get medical attention.
Do not eat, drink or smoke while using.
Remove protective clothing and wash hands and face thoroughly before meals and after work.
Wash protective clothing after work.
Avoid contamination of any water supply with chemical or empty container.

DIRECTIONS FOR USE:

MIXING:

Add the full quantity of the product to the partly filled spray tank whilst agitating. Fill tank and agitate to ensure thorough mixing.

APPLICATION:

AGPRO BIO-Safe™ may be applied through all conventional spray equipment including gun spraying, and knapsack spraying. Rainfall within 3 hours after application may reduce control.

APPLICATION RATES:

Rate	Critical Comment
70ml per 1 litre of water and spray to saturate weeds.	A second application after 3-4 days will assist control. Repeat treatments on undisturbed soil will reduce weed emergence.
On harder to kill weeds increase the rate to 100ml of per 1litre of water.	

RESISTANCE MANAGEMENT:

Not relevant to weeds being treated.

WITHHOLDING PERIOD:

Not applicable.

COMPATIBILITY:

Compatible with most commonly use herbicides.

SPRAY DRIFT WARNING:

Beware: Apply this product carefully. Spray drift may cause serious damage to other desirable plants. Ensure compliance with relevant regional air management plans.

SPILLAGE:

Wear appropriate protective clothing and prevent material from entering waterways. Absorb spills with inert material and place in waste containers. Wash area with water and absorb with further inert material. Dispose of waste safely (such as in a suitable landfill).

CONTAINER DISPOSAL:

Triple rinse container and add rinsate to spray tank. Burn in an appropriate incinerator, if circumstances such as wind direction permit. Otherwise crush or puncture and bury in a suitable landfill, or if appropriate, recycle.

SHELF LIFE:

When stored at temperatures between 8 - 28¼C, this product should show no significant degradation for two years from the date of manufacture. Contact us for further information about the use of product that is older than this.

CONDITIONS OF SALE:

The use of **AGPRO BIO-Safe™** being beyond the control of the manufacturer, no warranty expressed or implied is given by **AGPRO NZ LTD** regarding its suitability, fitness or efficiency for any purpose for which it is used by the buyer, whether in accordance with directions or not, and **AGPRO NZ LTD** accepts no responsibility for any consequence whatsoever resulting from the use of this product.