

READ THE DIRECTIONS FOR USE INFORMATION BEFORE USING THIS
PRODUCT

AGPRO DIURON 800

AGPROFLOWABLE HERBICIDE

For the control of a wide range of weeds in asparagus, grapes, kiwifruit, strawberries, gladioli, narcissus, tulips, iris, dry irrigation channels and drainage ditches.

Active Ingredient: 800g/litre diuron in the form of a suspension concentrate.

.NET CONTENTS: 20 Litres
Batch No:
Date of Manufacture:
HSNO Approval number:
Registered pursuant to the ACVM Act
1997 No. P4634
See www.nzfsa.govt.nz/acvm for
registration conditions.

Registered to and distributed by:
AGPRO NZ LTD
PO BOX 58-963, GREENMOUNT
10 POLARIS PLACE, EAST TAMAKI
AUCKLAND
PH 0-9-273 3456
FAX 0-9-273 3457
Web: www.agpro.co.nz

PRECAUTIONS:

Store in the original unopened container in a cool dry place, avoiding sunlight and away from foodstuffs and drink containers.

Do not get into eyes or on skin. In case of contact, flush skin or eyes with plenty of water for 15 minutes; for eyes get medical attention.

Do not eat, drink or smoke while using.

Remove protective clothing and wash hands and face thoroughly before meals and after work.

Wash protective clothing after work.

Avoid contamination of any water supply with chemical or empty container.

HAZARD CLASSIFICATIONS: 6.1D, 6.4A, 6.8B, 6.9A, 9.1A, 9.2A, 9.3C

FIRST AID:

- If swallowed do NOT induce vomiting.
- For advice contact National Poisons Centre (0800 POISON 764-766) or a doctor immediately.
- If skin or hair contact occurs, remove contaminated clothing and flush skin and hair with running water.
- If splashed in eyes, wash out immediately with water.

GENERAL INSTRUCTIONS

This product can be used for non-selective weed control on waste land, dry irrigation channels and drainage ditches. It can be used for the selective control in asparagus, grapes, strawberries, kiwifruit, gladioli, narcissus, tulips and iris. As AGPRO DIURON 800 is mainly absorbed through the roots of germinating weeds, it requires moisture to transport the product into the germinating zone of the weeds.

MIXING:

Invert container several times before opening. Stir well. Add the full quantity of product to the partly filled spray tank while agitating. Fill tank with water and agitate to ensure thorough mixing and continue agitation while spraying. Agitate vigorously from the bottom if allowed to stand. Reseal part used containers immediately.

APPLICATION:

Apply using conventional spray equipment. Care should be taken to avoid excessive spray overlap. Clean spray equipment after use. To be effective the chemical must be applied before weeds emerge and irrigation or light rain after application is necessary to allow the product to be moved to the root zone of the germinating weeds. Where dense growth is present add a suitable knockdown herbicide such as AGPRO Glyphosate 360.

COMPATIBILITY:

This product is compatible with all commonly used knockdown herbicides, however a test run is advisable.

SPRAY DRIFT WARNING:

Beware: Apply this product carefully. Spray drift may cause serious damage to other desirable plants. Ensure compliance with relevant regional air management plans.

CONTAINER DISPOSAL:

Triple rinse container and add rinsate to spray tank. Burn in an appropriate incinerator, if circumstances such as wind direction permit. Otherwise crush or puncture and bury in a suitable landfill, or if appropriate, recycle.

SHELF LIFE:

When stored at temperatures between 8 - 28¼C, this product should show no significant degradation for two years from the date of manufacture. Contact us for further information about the use of product that is older than this.

SPILLAGE:
Wear appropriate protective clothing and prevent material from entering waterways. Absorb spills with inert material and place in waste containers. Wash area with water and absorb with further inert material. Dispose of waste safely (such as in a suitable landfill).

SPILLAGE:

Wear appropriate protective clothing and prevent material from entering waterways. Absorb spills with inert material and place in waste containers. Wash area with water and absorb with further inert material. Dispose of waste safely (such as in a suitable landfill).

WITHHOLDING PERIOD: Not applicable

DIRECTIONS FOR USE**WEEDS CONTROLLED**

AGPRO DIURON 800 controls the following list of weeds at germination. It has little effect on established weeds. Annual poa, annual thistle, amaranthus spp, black nightshade, buttercup, chickweed, fathen, fleabane, fumitory, hemlock, henbit, mallows, redroot, scarlet pimpernel, spurrey, summer grass, stinking mayweed, shepherds

purse, twin cress, wild portulaca, wild turnip, willow weed and yarrow. It does not control Californian thistle, convolvulus, docks, Indian doab, nutgrass, oxalis, paspalum, plantains, sorrell and speedwells.

CROP	RATE/HA	CRITICAL COMMENT
Asparagus— established	1-2 litres/ha	On sandy or silty soils low in clay content or organic matter.
	2-4 litres/ha	On heavy clay soils or soils high in organic matter.
	In 250 - 500 litres water/ha	Apply prior to weed emergence and before the cutting season commences. If rainfall is expected a second application may be made immediately following harvest, however do not exceed 3.3 litres per season.
Bulbs-gladiolus, narcissus, tulip and iris	2 litres/ha in 250-500 litres in water/ha	Apply after the bulbs have been planted and before emergence of the tips. Do not apply post emergence or injury may result.
Grapes	2-4 litres/ha	On sandy or silty soils low in clay content or organic matter
	3-5 litres/ha	On heavy soils or soils high in organic matter.
	In 250-500 litres water /ha	Apply only to established vineyards at least 3 years old. Apply just prior to germination and growth of annual weeds in spring. Add AGPRO Glyphosate 360 for control of established weeds
Kiwifruit	Use same rate as for grapes	Apply just prior to germination and growth of annual weeds. Add AGPRO Glyphosate for control of established weeds.
Non selective vegetation control- non cropland	9-11litres/ha	For control of annual weeds
	11-36litres/ha in 250-500litres /ha water/ha.	For longer term control of annual and perennial weeds.
Waterways - drains, irrigation channels	21-24litres in 250-500litres water/ha.	Apply only when waterways are dry and at least 8 weeks before water is to be carried. Do not treat any area into which roots of trees or any desirable plants may extend.
Strawberries	2 litres/ha	Make one application per year between rows after transplanting before weeds emerge or before mulching. Apply only to strawberries grown in polythene. Do not make broadcast application.

CONDITIONS OF SALE:

The use of AGPRO DIURON 800 flowable herbicide being beyond the control of the manufacturer, no warranty expressed or implied is given by AGPRO NZ LTD regarding its suitability, fitness or efficiency for any purpose for which it is used by the buyer, whether in accordance with the directions or not and AGPRO NZ LTD accepts no responsibility for any consequence whatsoever resulting from the use of this product. For control of a wide range of weeds in asparagus, grapes, kiwifruit, strawberries, gladioli, narcissus, tulips, iris, dry irrigation channels and drainage ditches.