

Read the directions for use information before using this product.

AGPRO METURON Selective Herbicide

Active Ingredient 600g/kg of metsulfuron in the form of a water dispersible granule.

For the control of gorse and other scrub weeds in pasture, forest site preparation and waste areas.

NET CONTENTS: 500g
Batch No:
Date of Manufacture:
HSNO Approval number:
Registered pursuant to the ACVM Act 1997
No. P5306
See www.nzfsa.govt.nz/acvm for registration conditions.

Registered to and distributed by:
AGPRO NZ LTD
PO BOX 58-963, GREENMOUNT
10 POLARIS PLACE, EAST TAMAKI
AUCKLAND
PH 0-9-273 3456
FAX 0-9-273 3457
Web: www.agpro.co.nz

HAZARD CLASSIFICATIONS: 6.3B, 6.4A, 9.1A, 9.2A

WITHHOLDING PERIOD

Dairy cows should not be allowed into treated areas for at least three days after spraying. Other stock should be kept out of sprayed areas for at least one day.

RESISTANT WEEDS WARNING

AGPRO METURON is a member of the sulfonylurea group of herbicides. Some naturally occurring weed biotypes resistant to sulfonylureas may exist. These resistant weeds will not be controlled by AGPRO METURON or other sulfonylurea herbicides and may become dominant in that field. Cultural practices such as tillage, preventing weed escapes from going to seed and using herbicides with different modes of action within and between seasons can aid in delaying the proliferation and possible dominance of resistant weed biotypes.

MARKER DYES

AGPRO METURON is compatible with commonly used marker dyes but tank mixtures including marker dyes based on acetic acid should be used within 24 hours.

DECONTAMINATING EQUIPMENT

All spraying and mixing equipment should be thoroughly decontaminated after using AGPRO METURON as this product is a very effective herbicide in very low concentrations. Rinse all equipment and hoses thoroughly with clean water, (at least 10 minutes), then rinse with a 3% sodium hypochlorite solution. Then rinse all equipment and hoses with clean water again to remove the chlorine solution.

CONDITIONS OF SALE:

The use of AGPRO METURON being beyond the control of the manufacturer, no warranty expressed or implied is given by AGPRO NZ LTD regarding its suitability, fitness or efficiency for any purpose for which it is used by the buyer, whether in accordance with the directions or not and AGPRO NZ LTD accepts no responsibility for any consequence whatsoever resulting from the use of this product.

WEED CONTROL CHART - Application Rates

Weed	Handgun Rate/ 100L water	Knapsack Rate/ 10L water	Mistblower Rate/ 10L water	Aerial or Boom Rate/ Hectare	Critical Comments
Barberry	35g	5g	10g	-	Apply between November to May. Spray single bushes less than 3 m high for best results.
Blackberry	20-35g	5g	10g	200-300g	Apply between November to June. Best results are obtained when applied from January to May and prior to leaf discolouration. Use the higher rate in Northland and East Coast of North Island.
Bracken	35g	5g	10g	170g	Apply between January to June when fronds are unfurled, but before frost browning. Best results will be obtained when frond re-growth is sprayed after burning.
Broadleaf Weeds	5g	1g	2g	50g	Apply when the weeds are actively growing in drains, roadsides, industrial sites and waste areas.
Broom	35g	5g	10g	-	Apply from November to May to actively growing plants. Ensure AGPRO Organ silicone is added at 100mls/100litres.
Fennel	5g	1g	2g		Apply when weeds are actively growing.
Gorse	20g	5g	10g	0.5kg	Apply at any time of the year. For ULV spraying increase the rate to 20g/L. It is essential to add organ silicone at the following rates for all gorse spraying. Handgun: 100ml/100 litres. Knapsack: 10ml/100 litres. Aerial or ground boom: 400ml/100 litres.
Seedling gorse	-	-	-	170g	Apply at any time of year. Add AGPRO Organ silicone at 400ml/100 litres.
Hawthorn	35g	5g	10g	-	Apply from November to March. Spray single bushes less than 3 m high for best results.
Horehound	-	-	-	75g	Apply when weeds are actively growing.
Inkweed	25g	5g	10g	50g	Apply when weeds are actively growing.
Manuka	35g	5g	10g	-	Ensure complete spray coverage is obtained.
Matagouri	25g	5g	10g	170g	Apply December to April when actively growing.
Old Mans Beard	35g	5g	-	170g	Apply from November to May. Add AGPRO Organ silicone at 100mls/100 litres.
Privet	35g	5g	10g	-	Apply when actively growing during Spring or Autumn.
Ragwort /Thistle	35g	1g	-	-	Apply on ragwort from the rosette stage to late flowering, however for best results spray at the rosette stage. Spray into the centre of the plant. Use the higher rate on multicrown plants.
Sweet Brier	35g	5g	10g	170g	Apply from November to April. Ensure complete coverage is achieved.
Tauhinu	-	-	-	300g	Apply when actively growing. Spray single bushes less than 3m high for best results.
Tutu	25g	5g	10g	-	Apply when actively growing.
Wild Ginger	25g	5g	10g	-	Apply from Spring to late Autumn.
Willows	35g	5g	10g	-	Apply from November to April. Ensure complete coverage is achieved.