

WARNING

Keep out of reach of children

ECOTOXIC

AGPRO Picloram 20G

Contains 20g/kg picloram as the amine salt in the form of a granule.

For the control of blackberry, broom, gorse, ragwort, sweet brier, thistles, tutsan and other difficult to kill perennial weeds.

Read the directions for use before using this product.

Date of manufacture:

Registered pursuant to the ACVM Act 1997 No:P8133
See www.nzfa.govt/acvm for registration conditions.
Approved pursuant to HSNO Act 1996 No HSR000554
See www.ermanz.govt.nz for approval conditions

Net contents: 5kg & 25kg

Batch number:

Registered to and distributed by:

AGPRO NZ Ltd

10 Polaris Place,
East Tamaki
P O Box 58-963,
East Tamaki
Auckland.
Ph 09 273 3456
Fax 09 273 3457

Website: www.agpro.co.nz

HAZARD CLASSIFICATIONS: 6.5B, 9.1D, 9.2A.

WARNING:

Harmful: May cause sensitisation from prolonged skin contact.

Harmful to aquatic organisms. Selective herbicide - very toxic to some plant species.

PRECAUTIONS:

STORAGE: Keep out of reach of children. Store in original container, tightly closed and away from foodstuffs. Store in a cool dry well ventilated area, avoiding sunlight and temperatures below -4°C and above 35°C. When stored appropriately this product should show no significant degradation for

two years from the date of manufacture. Contact your supplier for information about the use of product that is older than this.

PERSONAL HYGIENE: Do not eat, drink or smoke while using. Remove protective clothing and wash hands and face thoroughly before meals and after work.

TRACKING: This product does not need to be tracked.

PRODUCT DISPOSAL: Dispose of this product only by using according to the label directions or if this is not possible alternatively contact your supplier for advice.

CONTAINER DISPOSAL: Dispose at an approved landfill. Do not burn. Do not use the container for any other purpose.

ENVIRONMENTAL: Use this product carefully. Do not contaminate aquatic environments with product or empty container.

First Aid: If swallowed DO NOT induce vomiting. If splashed in eyes, wash out immediately with water. If skin or hair contact occurs, remove contaminated clothing and flush with water. For advice call the National Poisons Centre 0800 POISON (0800 764 766) (24 hours) or a doctor immediately.

Emergency Response: In a transport emergency call 111, Police or Fire Brigade.

Spillage: Exclude non-essential people from the area. Wear protective clothing and prevent chemical from entering waterways.

Sweep up and place in a waste container. Dispose of safely to an approved landfill.

Safety Data Sheet: Additional information is provided in the safety data sheet. Phone 09 273 3456 or visit www.agpro.co.nz for a copy.

GENERAL INFORMATION

AGPRO Picloram 20G is a selective herbicide for spot application onto brushweeds and broadleaf weeds in areas where spraying is impractical or undesirable.

NOTE: The product is damaging to clovers. The treated area should be kept to a minimum and should not be used on peat or heavy clay soils.

DIRECTIONS FOR USE

APPLICATION:

AGPRO Picloram 20G should be applied over the top of the plant and in the case of brushweeds application should extend out to 300mm beyond the dripline. On steep slopes more granules should be applied on the uphill side as some downward leaching will occur. Best results are obtained under favourable growth conditions. Sufficient soil moisture must be present at the time of application to move the active ingredient into the soil. Application under dry conditions will give poor results.

WITHHOLDING PERIOD:

Poisonous plants may become more palatable after treatment, and livestock should be kept out until plants have died down. Otherwise no withholding period is required.

RATES OF APPLICATION:

Dock, nodding thistle and ragwort: Use 2g per plant (This equates to approx one level teaspoon and should be applied to the crushed centre of each plant).

Broom, fennel, goats rue, hemlock, inkweed, lupin, Mexican devil weed, St Johns wort and woolly nightshade: Use 30g per square metre. Best results are achieved on light sandy soils or stony soils with application being made in late winter to late spring.

Barberry (small bushes), blackberry, boxthorn (seedling), gorse, heath, Himalayan honeysuckle, horehound, matagouri, sweet brier, tutsan and tutu: Use 55g per square metre. Best results are achieved on light sandy soils or stony soils with application being made in late winter to late spring.

CONDITIONS OF SALE:

The use of AGPRO Picloram 20G herbicide being beyond the control of the manufacturer, no warranty expressed or implied, beyond that cover under the Fair Trading Act is given by AGPRO NZ Limited regarding its suitability, fitness or efficiency for any purpose for which it is used by the buyer, whether in accordance with the directions or not and AGPRO NZ Limited accepts no responsibility for any consequence whatsoever resulting from the use of this product.