

READ PRECAUTIONS BEFORE OPENING
READ THE DIRECTIONS FOR USE INFORMATION BEFORE USING THIS PRODUCT.

AGPRO **PROPAZINE 500** FOWABLE HERBICIDE

**For post-emergence control of selected weeds in carrots, parsnips,
and pine and douglas fir seed beds.**

Active Ingredient: 494g/litre propazine in the form of a suspension concentrate.

NET CONTENTS: Litres20

Batch No:

Date of Manufacture:

HSNO Approval number:

Registered pursuant to the ACVM Act

1997 No. P4608

See www.nzfsa.govt.nz/acvm for
registration conditions.

Registered to and distributed by:

AGPRO NZ LTD

PO BOX 58-963, GREENMOUNT

10 POLARIS PLACE, EAST TAMAKI

AUCKLAND

PH 0-9-273 3456

FAX 0-9-273 3457

Web: www.agpro.co.nz

PRECAUTIONS:

Store in the original unopened container in a cool dry place, avoiding sunlight and away from foodstuffs and drink containers.

Do not get into eyes or on skin. In case of contact, flush skin or eyes with plenty of water for 15 minutes; for eyes get medical attention.

Do not eat, drink or smoke while using.

Remove protective clothing and wash hands and face thoroughly before meals and after work.

Wash protective clothing after work.

Avoid contamination of any water supply with chemical or empty container.

HAZARD CLASSIFICATIONS: 6.1D, 6.3B, 6.4A, 6.7B, 9.1A, 9.2A

FIRST AID:

- If swallowed do NOT induce vomiting.
- For advice contact National Poisons Centre (0800 POISON 764-766) or a doctor immediately.
- If skin or hair contact occurs, remove contaminated clothing and flush skin and hair with running water.
- If splashed in eyes, wash out immediately with water.

GENERAL INSTRUCTIONS

This product is a pre-emergent herbicide which selectively controls certain broadleaf weeds and grasses in various crops. AGPRO PROPAZINE 500 is absorbed through the roots of germinating weeds and inhibits photosynthesis.

MIXING:

Invert container several times before opening. Stir well. Add the full quantity of product to the partly filled spray tank while agitating. Fill tank with water and agitate to ensure

thorough mixing and continue agitation while spraying. Agitate vigorously from the bottom if allowed to stand. Reseal part used containers immediately.

APPLICATION:

Apply using conventional spray equipment. Care should be taken to avoid excessive spray overlap. Clean spray equipment after use. To be effective the chemical must be moved to the root zone of the germinating weeds, thus irrigation or light rain after application is necessary for good results.

COMPATIBILITY:

This product is compatible with all commonly used agricultural insecticides, fungicides and herbicides, however a test run is advisable.

RESISTANCE MANAGEMENT:

Some annual weeds may be resistant. Use an alternative herbicide to remove them.

DRIFT WARNING:

Beware: Apply this product carefully. Spray drift may cause serious damage to other desirable plants. Ensure compliance with relevant regional air management plans. Aerial application: To minimise the risk of damage from spray drift, especially where susceptible vegetation is within 500 metres, use low drift-nozzles which minimise the production of small droplets (<100microns). Preferable to spray in a light constant breeze that is away from sensitive crops.

PROTECTION TO CROP, NATIVE AND NON-TARGET PLANTS:

When spraying for total weedkilling purposes, care should be taken not to spray beds where desirable annual plants are to be grown. Avoid drift of spray onto garden beds and plants.

SHELF LIFE:

When stored at temperatures between 8 - 28¼C, this product should show no significant degradation for two years from the date of manufacture. Contact us for further information about the use of any product that is older than this.

SPILLAGE:

Wear appropriate protective clothing and prevent material from entering waterways. Absorb spills with inert material and place in waste containers. Wash area with water and absorb with further inert material. Dispose of waste safely (such as in a suitable landfill).

CONTAINER DISPOSAL:

Triple rinse container and add rinsate to spray tank. Burn in an appropriate incinerator, if circumstances such as wind directions permit. Otherwise crush or puncture and bury in a suitable landfill, or if appropriate, recycle.

DIRECTIONS FOR USE

CROP	RATE/HA	CRITICAL COMMENT
Carrots & Parsnips	2-4L/ha in 500 L water.	Use the lower rate on sandy or silty soils and the higher rate on heavy clay soils or soils high in organic matter. Must be applied within 2-3 days of sowing and prior to emergence of the crop or weeds.
Radiata Pine seedbeds	2L/ha in 500L water	Apply within 2-3 days after sowing or 1 month after total emergence of the seedlings, ensuring that the seedbed has been freshly weeded.
Douglas Fir	2 kg/ha in 500L water	Apply at least 6 weeks after total emergence of the seedlings, ensuring that

seedbeds		the seedbed has been freshly weeded.
----------	--	--------------------------------------

CONDITIONS OF SALE:

The use of AGPRO PROPAZINE 500 flowable herbicide being beyond the control of the manufacturer, no warranty expressed or implied is given by AGPRO NZ LTD regarding its suitability, fitness or efficiency for any purpose for which it is used by the buyer, whether in accordance with the directions or not and AGPRO NZ LTD accepts no responsibility for any consequence whatsoever resulting from the use of this product