

KEEP OUT OF REACH OF CHILDREN  
CAUTION  
READ THE DIRECTIONS FOR USE BEFORE USING THIS PRODUCT.

# AGPRO TERMINATE FLOWABLE HERBICIDE

**For non-selective weed control in established crops of apples, citrus, grapes, kiwifruit, forestry and in non-cropland areas.**

Active Ingredient: 388g/litre terbuthylazine and 112g/litre glyphosate in the form of a suspension concentrate.

NET CONTENTS: 20 Litres  
Batch No:  
Date of Manufacture:  
HSNO Approval number:  
Registered pursuant to the ACVM Act  
1997 No. P5309  
See [www.nzfsa.govt.nz/acvm](http://www.nzfsa.govt.nz/acvm) for  
registration conditions.

Registered to and distributed by:  
AGPRO NZ LTD  
PO BOX 58-963, GREENMOUNT  
10 POLARIS PLACE, EAST TAMAKI  
AUCKLAND  
PH 0-9-273 3456  
FAX 0-9-273 3457  
Web: [www.agpro.co.nz](http://www.agpro.co.nz)

## **WARNING:**

This product may be harmful if swallowed, inhaled or absorbed through the skin.

## **PRECAUTIONS:**

Store in the original unopened container in a cool dry place, avoiding sunlight and away from foodstuffs and drink containers.

Do not get into eyes or on skin. In case of contact, flush skin or eyes with plenty of water for 15 minutes; for eyes get medical attention.

Do not eat, drink or smoke while using.

Remove protective clothing and wash hands and face thoroughly before meals and after work.

Wash protective clothing after work.

Avoid contamination of any water supply with chemical or empty container.

## **HAZARD CLASSIFICATIONS: 6.4A, 9.1A, 9.2A**

### **FIRST AID:**

- If swallowed do NOT induce vomiting.
- For advice contact National Poisons Centre (0800 POISON 764-766) or a doctor immediately.
- If skin or hair contact occurs, remove contaminated clothing and flush skin and hair with running water.
- If splashed in eyes, wash out immediately with water.

## **GENERAL INSTRUCTIONS**

AGPRO TERMINATE is a combination herbicide for post-emergent control of grasses and broadleaf weeds in established crops of apples, citrus, grapes, kiwifruit, forestry and non-cropland areas. This product is absorbed by foliar and root uptake. The

product works by inhibiting photosynthesis.

**MIXING:**

Invert container several times before opening. Stir well. Add the full quantity of product to the partly filled spray tank while agitating. Fill tank and agitate to ensure thorough mixing and continue agitation while spraying. Agitate vigorously from the bottom if allowed to stand.

**APPLICATION:**

AGPRO TERMINATE may be applied through all conventional spray equipment. Avoid any spray contact with crop foliage, fruit or green bark. In forestry apply as a pre-plant spray or if spraying after planting, ensure that the trees are well shielded from any spray drift. Store in a cool place and invert container three times before use. Do not use strainers finer than 300 micrometres. Do not store or mix AGPRO TERMINATE in non-galvanised or unlined steel containers.

**PROTECTION OF CROP, NATIVE AND NON-TARGET PLANTS:**

When used for total vegetation control purposes, care should be taken not to spray beds where desirable annual plants are to be grown. Avoid drift of spray on to garden beds and plants. Do not use in channels or drains or in other situations where roots of desirable plants or shrubs may extend. Do not use near newly planted shrubs, young ornamentals, and other species with surface roots.

**SPRAY DRIFT WARNING:**

**Beware: Apply this product carefully. Spray drift may cause serious damage to other desirable plants.**

It is an offence under the Pesticides Regulations to use this product in a manner which results in damage outside the treated property.

**DIRECTIONS FOR USE**

**WEEDS CONTROLLED**

AGPRO TERMINATE controls a wide range of grasses and broadleaf weeds. Visible effects may not be apparent for 7-14 days.

CROP	RATE	CRITICAL COMMENT
<b>Orchards</b> Apples, Citrus Grapes	6-10 litres/ha in 200-300 litres of water.	Use the lower rate on annual weeds and the higher rate on perennial weeds. More difficult to control weeds such as couch, kikuyu and paspalum will not be controlled. Do not use on trees or vines less than 3 years old. Do not allow the spray to contact any crop foliage, fruit or green bark.
Kiwifruit	6-10 litres /ha in 200-300 litres of water.	Use the rates as above. Do not use around vines less than 1 year old. Apply only when the vines are dormant. Do not allow the spray to contact any crop foliage, fruit or green bark.
<b>Forestry</b> Radiata pine, Douglas fir	12-25 litres/ha in 200-300 litres of water.	Pre-plant: Apply prior to planting, allowing sufficient time for the grasses to brown-off before tree planting. In cold weather, spots will take longer to brown-off. The lower rate will provide only short residual activity.
Eucalyptus nitens	12-20 litres/ha	
	Knapsack application. 1.5-2.5ml/square metre.	Post-plant: Shield the trees to prevent any spray drift contacting the trees.
<b>Non Cropland</b> Industrial areas, roadsides, footpaths, fencelines, etc.	15-25 litres/ha in 200-300 litres of water.	Apply sufficient spray to evenly cover the vegetation. Use the lower rate on light soils and the higher rate on soils with a high clay content and where long term weed control is desired.

***CONDITIONS OF SALE:***

The use of AGPRO TERMINATE flowable herbicide being beyond the control of the manufacturer, no warranty expressed or implied is given by AGPRO NZ LTD regarding its suitability, fitness or efficiency for any purpose which it is used by the buyer, whether in accordance with the directions or not and AGPRO NZ LTD accepts no responsibility for any consequence whatsoever resulting from the use of this product.